



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

the family possessing Property II is the set of conics having minor axes of an arbitrary constant length.

11. FOR THE LAW OF THE DIRECT DISTANCE. All orbits described under a force varying directly as the distance, are conics whose centers lie at the center of force. The force, expressed in terms of the constants of the conics, is given by

$$(24) \quad f = h^2 / a^2 b^2 u;$$

so that $h^2 / a^2 b^2$ must be constant for all the orbits. If also h is proportional to a/b^2 , it is necessary that b^6 , and hence also b , remain invariant. Thus, for this law, the family possessing Property I is the set of conics having minor axes of a given length. The constant of areas must vary as the major axis in this family. To have Property I at the pericenters, interchange b and a in the preceding statements. Each of these families has Property III also.

Finally, to have the product of the apsidal distances constant requires the constancy of $a.b$ and also of h . Hence, for the law of the direct distance, the family possessing Property II is the set of conics having the product of the axes constant, or what is equivalent for ellipses, having a given (arbitrary) area.

ON THE IRREDUCIBILITY OF CERTAIN POLYNOMIALS.

By JACOB WESTLUND, Purdue University.

The object of the following note is to determine whether the two polynomials

$$\begin{aligned} f_1(x) &= (x-a_1)(x-a_2) \dots (x-a_n) - 1 \text{ and} \\ f_2(x) &= (x-a_1)(x-a_2) \dots (x-a_n) + 1, \end{aligned}$$

where a_1, a_2, \dots, a_n are distinct integers, are reducible or irreducible.

Let us first consider $f_1(x)$. If $f_1(x)$ were reducible we would have

$$f_1(x) = \phi(x)\psi(x),$$

where $\phi(x)$ is irreducible and of a lower degree than n . Then since

$$\phi(a_i)\psi(a_i) = -1, \quad i=1, 2, \dots, n,$$

we must have

$$\phi(a_i) = \pm 1 \text{ and } \psi(a_i) = \mp 1.$$

Hence,

$$\phi(a_i) + \psi(a_i) = 0, \quad i=1, 2, \dots, n;$$

and hence the equation

$$\phi(x) + \psi(x) = 0,$$

whose degree is less than n , has n distinct roots, which is impossible. Hence, $f_1(x)$ is always irreducible.

Let us next consider $f_2(x)$. If $f_2(x)$ were reducible, we would have

$$f_2(x) = \phi(x)\psi(x),$$

where $\phi(x)$ is irreducible and of a lower degree than n . Then reasoning in the same way as in the first case we find that the equation

$$\phi(x) - \psi(x) = 0,$$

which is of a lower degree than n , has n distinct roots. But this is impossible, unless $\phi(x)$ and $\psi(x)$ are identically equal. Hence the only case when $f_2(x)$ is reducible is when it is a perfect square, in which case n of course must be even.



A METHOD OF COMPUTING LOGARITHMS.

By C. E. WHITE, Nashville, Tennessee.

From the expansion $f(x) = f(a) + (x-a)f'(a) + \frac{(x-a)^2}{2!}f''(a) + \dots$ we derive by letting $a = x+h$ and $a = x-h$,

$$f(x) = f(x+h) - hf'(x+h) + \frac{h^2}{2!}f''(x+h) - \frac{h^3}{3!}f'''(x+h) + \dots$$

$$f(x) = f(x-h) + hf'(x-h) + \frac{h^2}{2!}f''(x-h) + \frac{h^3}{3!}f'''(x-h) + \dots$$

The above expansions may be used to an advantage in computing log-